

Pediatric Pain Management (Non-Trauma)

WeightConversion Chart

KG	LB	KG	LB	KG	LB	KG	LB
1	2.20	31	68.34	61	134.48	91	200.62
2	4.41	32	70.55	62	136.69	92	202.83
3	6.61	33	72.75	63	138.89	93	205.03
4	8.82	34	74.96	64	141.10	94	207.23
5	11.02	35	77.16	65	143.30	95	209.44
6	13.23	36	79.37	66	145.50	96	211.64
7	15.43	37	81.57	67	147.71	97	213.85
8	17.64	38	83.78	68	149.91	98	216.05
9	19.84	39	85.98	69	152.12	99	218.26
10	22.05	40	88.18	70	154.32	100	220.46
11	24.25	41	90.39	71	156.53	101	222.67
12	26.46	42	92.59	72	158.73	102	224.87
13	28.66	43	94.80	73	160.94	103	227.08
14	30.86	44	97.00	74	163.14	104	229.28
15	33.07	45	99.21	75	165.35	105	231.49
16	35.27	46	101.41	76	167.55	106	233.69
17	37.48	47	103.62	77	169.76	107	235.89
18	39.68	48	105.82	78	171.96	108	238.10
19	41.89	49	108.03	79	174.16	109	240.30
20	44.09	50	110.23	80	176.37	110	242.51
21	46.30	51	112.44	81	178.57	111	244.71
22	48.50	52	114.64	82	180.78	112	246.92
23	50.71	53	116.84	83	182.98	113	249.12
24	52.91	54	119.05	84	185.19	114	251.33
25	55.12	55	121.25	85	187.39	115	253.53
26	57.32	56	123.46	86	189.60	116	255.74
27	59.52	57	125.66	87	191.80	117	257.94
28	61.73	58	127.87	88	194.01	118	260.15
29	63.93	59	130.07	89	196.21	119	262.35
30	66.14	60	132.28	90	198.42	120	264.55

Routine Medical Care
(Including pain scale)

- Evaluate mechanism of injury
- Assess the need for spinal precautions per **Spinal Motion Restriction** protocol

Attempt comfort therapies first:
Record level of pain by either asking patient to rate on scale 1-10 or using **FACES** scale

Administer one of the following for moderate to severe (4-10) pain.

- **Morphine Sulfate** 0.05mg/kg IV (< 1 year old)
0.1mg/kg IV (≥ 1 year old)
- **Ketorolac** 0.5 mg/kg IV/IO/IM (max 15 mg)
- **Acetaminophen** 15 mg/kg IV infusion over 15 minutes ¹*For pediatric patients weighing >50kg or 13 years old-refer to adult dose.

Reassess pain level 5-10 min after treatment

If pain remains severe (score: 8-10) **contact Medical Control** for additional orders

Have **Naloxone** readily available to treat narcotic-induced respiratory depression.



Wong-Baker FACES of pain rating scale.

¹DO NOT run IV Infusion faster than over 15 minutes, the patient may experience burning sensation. Acetaminophen IV has peak effect within 30 min and last 4-6 hours.